

FLTAX (5-FU/I-LV/PTX) 療法

| | | | |
|----|----|------------|----|
| 適応 | 胃癌 | 催吐性 リスク | 軽度 |
|----|----|------------|----|

| | | | Day1 | Day8 | Day15 | Day22 | Day28 |
|------|----------------------|------------|------|------|-------|-------|-------|
| PTX | 60mg/m ² | DIV, 60min | ● | ● | ● | | |
| I-LV | 250mg/m ² | DIV, 60min | ● | ● | ● | | |
| 5-FU | 500mg/m ² | IV, bolus | ● | ● | ● | | |

4週間毎, PDまで

| Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
|--------------------------------|--------|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| ① 生理食塩液注 | 50mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| デキサート注射液 | 8.25mg | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| ファモチジン注 | 20mg | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| DIV | 10min | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ② グラニセトロンバッグ3mg | 50mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| DIV | 30min | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ③ パクリタキセル 60mg/m ² | | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| 生理食塩液注 | 250mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| DIV | 60min | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ④ レボホリナート 250mg/m ² | | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| 生理食塩液注 | 250mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| DIV | 120min | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ⑤ 5-FU 500mg/m ² | | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | |
| IV | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

【前投薬】 レスタミンコーワ錠 50mg/経口