

XELIRI+Bev 療法

適応

大腸癌

催吐性
リスク

中等度

			Day1	Day8	Day15	Day21	
Capecitabine	800mg/m ² /回	1日2回, 経口	————→			14日間	
Bevacizumab	7.5mg/kg	DIV, 90→60→30min	●				
CPT-11	200mg/m ²	DIV, 120min	●				

3週間毎,

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① カペシタピン 800mg/m ² ×2		○	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	○													
経口		夕														朝													
② アロキシバッグ0.75mg 50mL		●																											
デキサート注射液 9.9mg		●																											
DIV 30min																													
③ アバスチン 7.5mg/kg		●																											
生理食塩液注 100mL		●																											
DIV 90→60→30min																													
④ イリノテカン 200mg/m ²		●																											
生理食塩液注 500mL		●																											
DIV 120min																													

【Capecitabine投与量】

体表面積	1回用量
1.31m ² 未満	900mg/回
1.31m ² 以上1.69m ² 未満	1,200mg/回
1.69m ² 以上2.07m ² 未満	1,500mg/回
2.07m ² 以上	1,800mg/回