

biweeklyPTX+S-1 療法

適応

胃癌

催吐性
リスク

軽度

			Day1	Day8	Day15	Day22	Day28
S-1	40mg/m ² /回	1日2回, 経口	→ 14日間				
PTX	120mg/m ²	DIV, 60min	●		●		

4週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① S-1 40mg/m ² ×2		○	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	○													
経口		夕														朝													
② 生理食塩液注 50mL		●														●													
デキサート注射液 8.25mg		●														●													
ファモチジン注 20mg		●														●													
DIV 15min																													
③ グラニセトロンバッグ3mg 50mL		●														●													
DIV 30min																													
④ パクリタキセル 120mg/m ²		●														●													
生理食塩液注 500mL		●														●													
DIV 120min																													

【前投薬】 レスタミンコーワ錠 50mg/経口

【S-1投与量】

体表面積	S-1初回基準量
1.25m ² 未満	40mg/回
1.25m ² 以上1.5m ² 未満	50mg/回
1.5m ² 以上	60mg/回