

SOX+Trastuzumab 療法

適応	胃癌	催吐性 リスク	中等度
----	----	------------	-----

			Day1	Day8	Day15	Day21
S-1	40mg/m ² /回	1日2回, 経口	—————→			14日間
L-OHP	130mg/m ²	DIV, 120min	●			
Trastuzumab	8mg→6mg/kg	DIV, 90→30min	●			

3週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① S-1 40mg/m ² ×2		○	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	○													
経口		夕														朝													
② アロキシバグ0.75mg 50mL		●																											
デキサート注射液 9.9mg		●																											
DIV 15min																													
③ トラスツズマブ 8mg→6mg/kg		●																											
蒸留水		●																											
生理食塩液注 250mL		●																											
DIV 90→30min																													
④ オキサリプラチン 130mg/m ²		●																											
5%ブドウ糖液 500mL		●																											
DIV 120min																													

【S-1投与量】

体表面積	S-1初回基準量
1.25m ² 未満	40mg/回
1.25m ² 以上1.5m ² 未満	50mg/回
1.5m ² 以上	60mg/回