

IRIS+Bev 療法

適応

大腸癌

催吐性
リスク

中等度

			Day1	Day8	Day15	Day22	Day28
S-1	40mg/m ² /回	1日2回, 経口	→ 14日間				
Bevacizumab	5mg/kg	DIV, 90→60→30min	●		●		
CPT-11	100mg/m ²	DIV, 120min	●		●		

4週間毎,

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
① S-1 40mg/m ² ×2		○	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	○														
経口		夕														朝														
② アロキシバグ0.75mg 50mL		●														●														
デキサート注射液 9.9mg		●														●														
DIV 15min																														
③ アバスチン 5mg/kg		●														●														
生理食塩液注 100mL		●														●														
DIV 90→60→30min																														
④ イリノテカン 100mg/m ²		●														●														
生理食塩液注 500mL		●														●														
DIV 120min																														

【S-1投与量】

体表面積

S-1初回基準量

1.25m²未満

40mg/回

1.25m²以上1.5m²未満

50mg/回

1.5m²以上

60mg/回