

weekly PAC 療法

適応

非小細胞肺癌

催吐性
リスク

軽度

| | | | Day1 | Day8 | Day15 | Day22 | Day28 |
|-----|---------------------|------------|------|------|-------|-------|-------|
| PTX | 70mg/m ² | DIV, 60min | ● | ● | ● | | |

4週間毎, PDまで

| Day | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | |
|-------------------------------|-----------|---|-----------|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|--|
| ① 生理食塩液注 | 50mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | | |
| デキサート注射液 | 6.6mg | ● | 初回は8.25mg | | | | | | ● | | | | | | | | ● | | | | | | | | | | | | | |
| ファモチジン注 | 20mg | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | | |
| | DIV 15min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ② グラニセトロンバッグ3mg | 50mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | | |
| | DIV 30min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ③ パクリタキセル 70mg/m ² | | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | | |
| 生理食塩液注 | 250mL | ● | | | | | | | ● | | | | | | | ● | | | | | | | | | | | | | | |
| | DIV 60min | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

【PTX前投薬】 レスタミンコーワ錠 50mg/経口