

Gefitinib/CBDCA+PEM 療法

適応

非小細胞肺癌

催吐性
リスク

中等度*

			Day1	Day8	Day15	Day21
Gefitinib	250mg/回	1日1回, 経口	—————→			
CBDCA	AUC 5	DIV, 60min	●			
PEM	500mg/m ²	DIV, 10min	●			

3週間毎, 4コース

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① イレッサ 250mg/回×1		◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎	◎							
経口																													
② グラニセトロンバッグ3mg 50mL		●																											
デキサート注射液 4.95mg		●																											
DIV 20min																													
③ アリムタ 500mg/m ²		●																											
生理食塩液注 100mL		●																											
DIV 10min																													
④ カルボプラチン AUC 5		●																											
5%ブドウ糖液 250mL		●																											
DIV 60min																													

【初回投与7日前から開始】パンビタン末：1.0g/回, 経口、連日

【初回投与7日前】メチコバル注：1,000μg/回、筋注, 9週毎

※最終投与日から22日目まで継続