

Trastuzumab+S-1 療法

適応

乳癌

催吐性
リスク

軽度

			Day1	Day8	Day15	Day22	Day29	Day36	Day42	
S-1	40mg/m ² /回	1日2回, 経口	→ 28日間							
Trastuzumab	4mg→2mg/kg	DIV, 90→30min	●	●	●	●	●	●		

6週間毎, PDまで

Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
① S-1 40mg/m ² ×2		○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
経口		夕																											
② トラスツズマブ 4mg→2mg/kg		●							●							●							●						
蒸留水		●							●							●							●						
生理食塩液注 250mL		●							●							●							●						
DIV 90→30min																													

Day	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56
① S-1 40mg/m ² ×2	○																											
経口	朝																											
② トラスツズマブ 4mg→2mg/kg	●							●																				
蒸留水	●							●																				
生理食塩液注 250mL	●							●																				
DIV 90→30min																												

【S-1投与量】

体表面積	S-1初回基準量
1.25m ² 未満	40mg/回
1.25m ² 以上1.5m ² 未満	50mg/回
1.5m ² 以上	60mg/回